

Cost / Carbon footprint / Efficiency evaluation for the treatment of TPH contaminated soils: off-site disposal vs. on site Bio- or Rhizo-degradation

Intersol 2019 – Lille, France – March 27th, 2019


Florian Le Chevalier, Romain Carthagena, Anouk Barrere,
Nathalie Montigny, Julie Estival, Sébastien Kaskassian

Context

Soils contaminated by drilling mud/slurry from oil extraction wells

- Mud / slurry fluidized with oil
 - diesel-like product, mainly C13-C25, 70% aliphatic HC
 - Average conc. ~ 30,000 mg/kg DW
- Site configuration:
 - Limited space, rural zone, no access to energy
 - Land to be restored by 2026
 - 16,000 m³ soil/sludge stock

→ Treatment options?



Comparison metrics of treatment options

How to evaluate cost – duration – efficiency?

- Off site solutions: Thermal Desorption vs. Biological

Cost – duration – efficiency = given by treatment facilities

- On site solutions: Thermal Desorption vs. Biological with / without vegetation growth

Cost – duration – efficiency = given by contractor for TD ... but for biological ???

→ Field scale feasibility study (biopiles) to better evaluate actual design parameters and treatment conditions

- **Biodegradation** (aerobic) vs. **Biodegradation enhanced by vegetation**
- Monitoring for ~2 years period
- Performance criteria: reduction in TPH conc. & toxicity, cost / duration / carbon footprint



Configuration of the field pilots (~100m³ each)

based on previous lab. exp. performed by MicroHumus

Biopiles “A” : biodegradation (21 months)



- Aeration by mechanical mixing every 2-3 months
- Under cover (rain-evaporation & light preservation)

Rhizopiles “C” : bio- (4 months) + vegetation (17 months)



- 0-4 months: same as Biopiles “A”
- @ 4, 10 and 15 months: (re)seeding Ray-grass and Alfalfa
- >4months : undisturbed and uncovered

*arable soil planted @2 months

Mixture (%DW)	A&C 4/6	A&C 4/6a	A&C 2/8	A&C 2/8a	Polluted Control	Arable soil*
Soil/sludge	64,9%	64,8%	84,3%	84,1%	100%	
Arable soil	31,2%	31,1%	13,2%	13,1%		100%
Compost	3,8%	3,8%	2,5%	2,5%		
Limestone	-	0,26%	-	0,25%		
Urea	0,02%	0,02%	0,02%	0,02%		

Compost	proportion
Pine bark 10/25 mm	70 %
Green vegetation 0/10 mm	30 %



Monitoring program and partnership

Compare all treatment conditions: sampling every 2-3 months for 21 months



- **In-situ measurement** in piles: pH, Temperature, and biogas production (CO₂-O₂-CH₄) → [Tauw France](#)
- **TPH concentrations** in replicate samples, distribution with depth in the piles (0-30 / 30-60 / 60-100 cm),
- **General soil quality**: metals, N/P/K balance and speciation, cationic exchange capacity, alkalinity, TOC, → [Synlab](#)
- **Microbial analysis**: RISA on Bacteria and Fungi, qPCR on total bacteria and total Fungi, qPCR on functional genes (TPH degraders) → [Enoveo](#)
- **Nematodes** index and diversity (NF/ISO 23611-4) → [Elisol](#)
- **Plants development**: sampling after plant growth
 - Plant growth (root and stem), biomass production → [Tauw France](#)
 - Omega 3 analysis in leaves (XP X31-233) → [LEB-Aquitaine](#)
 - TPH analysis in the biomass (plant uptake and accumulation) → [Eurofins](#)
- **TPH advance analysis & availability in soil matrices**:
 - in-depth chromatography, thermodesorption lab tests → [Univ. Lorraine \(LIEC/Géoressource\)](#)



Performance monitoring: TPH results

Biopiles “A” : biodegradation (8 months)

Rhizopiles “C” : bio- (4 months) + vegetation (4 months)







Is it biologically active or is there “something else” ? Do plants help ? Any matrix effect ?



Extrapolation for « complete degradation »

Theoretical treatment target = 1 500 mg/kg DW

Estimated duration (in months) of treatment based on 10 months monitoring (only)




- **Expected treatment duration?**
 - Biopiles: no significant differences
 - Rhizo-piles: same except for 1 pile
- **Enhancement by vegetation?**
 - Longer expected treatment duration with vegetation (w/o mechanical mixing): +45% to + 60%
 - No mixing & Heterogeneity in soils
- **Reduction in Toxicity (restore agriculture use)?**
- **What is the achievable residual concentration? → all TPH fractions 100% degradable?**




Indicators of biodegradation in the piles (after 10 months)

In-situ gas monitoring

BioPiles A



RhizoPiles C



In-situ gas

- O₂ uptake and CO₂ production = higher efficiency in Biopiles
- Rather stable pH

Good aeration relies on mechanical mixing (biopiles A) or root development (rhizopiles C)

Biomarkers


- Increase in total bacteria : higher in amended biopiles
- Evolution in specific genes (degraders for C5-C20 TPHs)
- Stable fungi and nematode populations

Bacterial population difference in Vegetated piles?


Chemical indicators as other proof of biodegradation processes?

Biomarkers monitoring

BioPiles




Amended BioPiles




Vegetation development & Ecotoxicity in Rhizopiles C (seeding in April-18)


Density & height after 2 months (June-18)



Omega3 index after 4 months (Aug.-18)



Biomass production after 6 months (Oct.-18)
















- Vegetation growth
 - arable soil = reference for plant growth
 - lower Alfalfa density & Omega 3 as well as total biomass production with high TPH or limestone content
- Other ecotoxicity tests (earthworm, daphnia, Fischeri) = no difference in piles / arable soil / raw contaminated soil
- No TPH measured in biomass → **possibility to use soils as agricultural land after treatment?**



Carbon footprint of remediation options







Carbon footprint computations

- REC model (Haselhoff et al., 2011)
- Dynamic spreadsheet for remediation options
- Inputs
 - Power source / nature (7 choices)
 - Consumption & fuels
 - Materials (pipes, sheets ...)
 - Reactions (chemical / biological)
 - Pollutants and expected efficiency
 - ...
- Output = CO2 emission
 - Equipement
 - Transport, vehicle type
 - Maintenance & monitoring
 - Raw material, wastes, effluents
 - ...

A	B	C	D	E	F	G	H	I	J	K	L	M
					Calculation model CO₂ for soil remediations <small>version 1.3.3-NL</small>							
												
					<u>Import screen</u> Enter the projectdata and click on the buttons to take out the import fields for the concerning component. Follow the questions and options for the concerning component and enter the correct data.							
					Projectname <<projectname>>							
					Remediationvariant <<remediationvariant>>							
					Date <<date>>							
					Model filled by <<name>>							
					<hr/>							
					 Excavation landsoil							
					 Groundwater extraction							
					 Groundwater purification							
					 In situ remediation Airsparging and Soil Vapor Extraction (SVE)							
					 In situ remediation Multi Phase Extraction (MPE)							
					 In situ remediation ISCO							
					 In situ remediation Biostimulation							
					 In situ remediation Thermal							
					 Monitoring and aftercare							
												
												
					<hr/>							
					Manual Import Output Datasheet 1 Datasheet 2 Datasheet 3 Calculations 							



Carbon footprint results



Overview of treatment alternatives for 16 000 m³

	Treatment	Time	Cost (€)	Carbon footprint (tons of CO ₂)	Remarks (+ / -)
Off site treatment	Thermal desorption	4 months*	3,8 M€	11 400 T CO ₂ 55% due to transport, 44% due to treatment	+ Residual concentration < 50 mg/kg - Transport outside France (ship) - Loss in soil functions
	Biological	4 months*	2,8 M€	6 300 T CO ₂ 98% due to transport	- Residual concentration # 500 mg/kg - Transport outside France (ship) +/- Increased value of soils?
On-site treatment	Biological	5 months* + 2-3 years**	2,4 M€	174 T CO ₂ 56% due to excavation, 32% due to treatment	- Residual concentration # 1500 mg/kg - Treatment time x mechanical aeration +/- New use for treated soils?
	Biological & Rhizo-dégradation	5 months* + 3-5 years**	2,1 M€	170 T CO ₂ 56% due to excavation, 32% due to treatment	- Residual concentration # 1500 mg/kg + Soil aeration due to roots growth - Longer treatment time + New use for treated soils: it seems so! + CO ₂ captation by biomass growth (~3 T)
	Thermal desorption	4 months* + 2 months**	2,8 M€	5 200 T CO ₂ 98% due to treatment	+ Residual concentration < 50 mg/kg - Losses in soil functions +/- Treated soil buried on site (excavation open pit)

* Duration for excavation & preparation works and soil loading (time for off site treatment not included).

** Duration of on-site treatment to achieve reduction in TPH concentrations (extrapolation from results gathered after 10 months monitoring)



Conclusions and pending work

CO₂ emission criterion = insight in Cost-Benefit Analysis for remediation options

- Shorter Time: Off-site treatment options save 2 to 5 years
- Lowest Costs: On-site Bio- / Rhizo- Degradation save > 1 M €
- Lowest Carbon footprint: On-site Bio- / Rhizo- Degradation save 5 to 10 K Tons CO₂
- Higher Efficiency: Thermal Desorption (Off- / On-Site) achieves residual < 1% initial

Questions to be addressed during remaining monitoring period (until Sept.)

- Design parameters & best treatment conditions
 - Thorough statistical, correlation & tendency analysis
- Residual effects (Toxicity) and soil use / valuation (agricultural use)
 - Other Ecotoxicity tests (functional responses & TPH transfer in agricultural crop)?
 - Public / Authority acceptance?
- Lowest achievable residual concentrations with on-site bio- or rhizo-degradation?
 - Bio-availability evaluation: thermal lab. essays (on-going work with GISFI-LIEC)



Thank you!



Sébastien Kaskassian



+33 (0)6 32 15 14 36



s.kaskassian@tauw.com



www.tauw.fr

