



# Study of contamination and toxicity of PFAS in drinking water in the Fos-Berre area and analytical developments for atmospheric biomonitoring using lichens

*M. Dijoux<sup>1</sup>, M. Denton<sup>2</sup>, A. Piram, S. Augy, A. Austruy, C. Loquais, A. Souloumiac, L. De Jong, E. Gonzales-Camoin, P-H. Villard, L. Terlier, P. Bremond, K. Benbrahim, J. Dron, P. Chamaret, X. Moreau, S. Gori, P. Wong–Wah–Chung, P. Doumenq*

<sup>1</sup>ABO-ERG Environnement, LCE (Aix-Marseille University)

<sup>2</sup>IMBE (Aix-Marseille University)



## Per- and polyfluoroalkylated substances (PFAS)



More than 12 000 molecules



Anthropogenic origin




Synthesized since 1940'

### Physico-chemical properties [1]

- Hydrophobicity
- Lipophobicity
- Surfactant
- Anti-adhesive
- Heat resistance and stability
- Chemical stability

### Toxicity [2, 3]



Thyroid  
Cholesterol  
Liver  
Cancers  
Endocrine disruptors  
Fertility & pregnancy

Many  
associated  
pathologies

November 2023 : 2 PFAS classified as **carcinogenic** and **possibly carcinogenic** to human [4]

### Increasing regulations




Drinking water (16/12/2020) [5]

Effluents (20/06/2023) [6]

Atmospheric emissions (31/10/2024) [7]

## PFAS exposure pathways



# MATISSE Project : Multi-scale PFAS contamination Study in industrial areas

- 01/2024-12/2026
- Part of the project France 2030
- Financed by Amidex foundation and ERG foundation
- Budget: 1,43 million €



## Occurrence

- Study of targeted PFAS in drinking water, groundwater and lichens
- Suspected analysis




## Multi-matrix transfers

- Soil, air, groundwater
- Comparison of a « manufacturing » site (Lyon) and a « user » site (Fos-Berre-Marseille)



## Eco- and human toxicity

- Invertebrate and human cell exposure to mixtures of PFAS



[8]



# Lichen biomonitoring

**Aim****Evaluation of the atmospheric exposition (NF X43-904)**

## Advantages of the lichens

- Lack of root system → Representative of atmospheric contamination → **Bioindicator**
- Stationary → **Spatial resolution**
- Integration of the data over several months → **Integrative bio-sampler**
- Can live in urban, industrial, rural and extreme areas (deserts, Antartica) and on all types of substrate (trees, rocks, buildings) → **Ubiquitous organisms**
- Already studied for metals, pesticides, PCBs, PAHs, PCDD-F

# Lichen biomonitoring



Techniques	PFAS in lichens	PFAS in other plant matrices <a href="#">[11]</a>	Studied parameters
Extraction method	Solid/liquid extraction	Simple agitation, ultra-sonication, QuEChERS, assisted solvent extraction	Orbital agitation Ultra-sonication
Extraction solvent	Methanol	Methanol, acetonitrile, MTBE, ethyl acetate or a mixture of solvents, sometimes with formic acid, ammonium acetate or ammonium hydroxide	Methanol, with and without formic acid or ammonium hydroxide
Purification	Sampli Q carbon <a href="#">[9]</a> Activated carbon <a href="#">[10]</a>	Carbon (solid phase extraction or dispersive (d)SPE) → graphitic carbon (ENVI-Carb™), carbon, C <sub>18</sub> , Ionic exchange (WAX)	Captiva EMR PFAS Food I ( <i>Agilent</i> ) ENVI-Carb™ ( <i>Supelco</i> ®)

## Lichen biomonitoring



Techniques	PFAS in lichens	PFAS in other plant matrices [11]	Studied parameters
Extraction method	Solid/liquid extraction	Simple agitation, ultra-sonication, QuEChERS, assisted solvent extraction	Orbital agitation Ultra-sonication
Extraction solvent	Methanol	Methanol, acetonitrile, MTBE, ethyl acetate or a mixture of solvents, sometimes with formic acid, ammonium acetate or ammonium hydroxide	Methanol, with and without formic acid or ammonium hydroxide
Purification	Sampli Q carbon [9] Activated carbon [10]	Carbon (solid phase extraction or dispersive (d)SPE) → graphitic carbon (ENVI-Carb™), carbon, C <sub>18</sub> , Ionic exchange (WAX)	Captiva EMR PFAS Food I (Agilent) ENVI-Carb™ (Supelco®)


## Lichen biomonitoring




Techniques	PFAS in lichens	PFAS in other plant matrices [11]	Studied parameters
Extraction method	Solid/liquid extraction	Simple agitation, ultra-sonication, QuEChERS, assisted solvent extraction	Orbital agitation Ultra-sonication
Extraction solvent	Methanol	Methanol, acetonitrile, MTBE, ethyl acetate or a mixture of solvents, sometimes with formic acid, ammonium acetate or ammonium hydroxide	Methanol, with and without formic acid or ammonium hydroxide
Purification	Sampli Q carbon [9] Activated carbon [10]	Carbon (solid phase extraction or dispersive (d)SPE) → graphitic carbon (ENVI-Carb™), carbon, C <sub>18</sub> , Ionic exchange (WAX)	Captiva EMR PFAS Food I (Agilent) ENVI-Carb™ (Supelco®)




## Extraction



## Extraction



## Purification



## Analysis



- Method developed for 49 PFAS (PFCAs, PFSAAs, Gen-X, ADONA, FTS, diPAPs, FASAs)
- LC-MS/MS method not sensitive enough for volatile PFAS (FTOHs)

LC-QqTrap 7500 (Sciex)

## Conclusion → Comparison of extraction recoveries and matrix effects

### Purification:

- EMR Captiva Food I → Ionic PFAS retained on the cartridge and could not be eluted
- ENVI-Carb™ → Good purification recoveries (70-120%)
  - Extracts were further diluted to reduce matrix effect

### Extraction solvent:

- MeOH 0,1% NH<sub>4</sub>OH → Best extraction recovery
- MeOH 0,1% FoA
- MeOH

### Extraction technique:

- Orbital agitation
- Ultra-sonication → Good extraction recoveries (70-120%), better repeatability and lower matrix effects

**Limit of quantification of the method : < 1 ng.g<sup>-1</sup> (dw) for most PFAS**

**Method to be validated and applied to a sampling campaign**

## Drinking water campaign



Sampling carried out with the help of the participatory science network of the IECP

- **28 tap waters**

from the Crau groundwater table or the Durance river

- **2 groundwaters**

private wells from the Crau groundwater table

- **1 field blank per sampling site** (MilliQ water)



Background map sources : ESRI

## Sampling and analytical method

**Analytical challenges:** contamination, loss, degradation [12]

→ **Adaptation of the protocols**



### Sampling

- HDPE bottles

### Conservation

- Addition of a preservative agent
- Frozen (-20 °C)

### Preparation

- Dilution in 60% MeOH [13]

### LC-MS/MS Analysis

- Within 2 weeks
- Adaptation of analytical chain

- Water-repellent clothes
- Food packagings
- Fluoropolymers
- PFAS containing chemicals



**35 PFAS**

Sciex Qtrap 7500

## Evaluation of the method accuracy:

- Water Certified Reference Material (CRM IRMM-428)

- All results within 30%  
of the certified values




- 5 duplicated samples analysed by a certified laboratory (Eurofins)

- 90% of the results within 30%  
of the certified values




## PFAS investigation in drinking water

Sum of PFAS concentrations in tap water (ng.L<sup>-1</sup>)





## PFAS investigation in drinking water



➔ Higher frequency of quantification for short-chain PFAS (3-8 carbons)


- Higher solubility in water
- Used in replacement of long-chain PFAS
- Degradation products of long-chain PFAS precursors

### Main uses of the PFAS found in the samples



## Comparison to the European regulation

Sum of PFAS concentrations in tap water (ng.L<sup>-1</sup>)



2020/2184 European Drinking Water

Directive (16/12/2020)

$\Sigma$  20 PFAS: 100 ng.L<sup>-1</sup>

Total PFAS: 500 ng.L<sup>-1</sup>


→ 10 of the quantified PFAS  
are part of the 20 regulated  
PFAS in drinking water

→ 1 site above the European  
regulation

## Alarming results ? A regulation-dependant interpretation

Total number of samples = 30

Number of samples exceeding the regulation





New French recommandation :  $\sum 4 \text{ PFAS (PFOA + PFOS + PFNA + PFHxS)} < 20 \text{ ng.L}^{-1}$  [14]

➔ The same sample exceeds the recommandation

# Conclusions for drinking water analysis

- ➔ Concentrations impacted by the location of the water catchment, explaining the **variability** in PFAS contamination **within** and **between** towns



Background map sources : ESRI

- ➔ Higher contamination for water catchments located downstream of the groundwater table, except for individual catchments
- ➔ Contamination is not higher in individual catchments than in collective catchments (despite additional treatments)
- ➔ Difference in contamination between neighbouring catchments is not yet explained

## Selecting PFAS for the study of toxicological and ecotoxicological properties



⇒ Based on

Occurrence

Abundance

Presence or absence  
of existing literature

Ability to recreate  
the mixtures

### Final choices

#### ❖ 3 PFAS for toxicological assays :

PFOS, PFHxA and PFPeA

#### ❖ 9 PFAS for ecotoxicological assays :

PFODA, PFOS, PFOA, PFHpA, PFHxS, PFHxA, PFBS, PFPeA, PFBA

## Toxicological assay



### Aim

Evaluate the potential effects of an exposure to PFAS for 24 h via the digestive tract

### Model

Caco-2 cell line (enterocyte phenotype)

### Technique

RT-qPCR



### Exposure concentrations

- ✓ PFOS : 7.7 ng/L
- ✓ PFHxA : 0.35 ng/L
- ✓ PFPeA : 90 ng/L
- ✓ **Mix of the 3**



### Genes of interest

- **Inflammation**  
(IL1b, IL6, TNFa)
- **Detoxification**  
(CYP1A, AhR)
- **Oxidative stress**  
(CAT, SOD2, GPx)
- **Epithelium integrity**  
(CLDN1 & CLDN 3)



Caco2 cells


Photo : M. Denton



## Toxicological assay : Preliminary results



### Relative expression of SOD2 (superoxide dismutase) in Caco2



→ No significant differences compared to solvent controls ( $P > 0.05$ )

→ Increase the number of replicates to strengthen statistical power

## Ecotoxicological assays

### Advantages of using *Hydra vulgaris* as a model

- **Invertebrate** (3R → replacing vertebrate models)
- **Diploblastic organism** → full exposure to the aquatic medium
- **Ability to regenerate** → similar to a hydra embryo, indication of the potential teratogenicity of a substance [15, 16]
- **Easy to rear** in a laboratory
- **Sequenced genome**
- **Multi-scale biological responses** : molecular / organism / population



## Ecotoxicological assays



Morphology and  
teratogenicity  
studies



Gene expression  
levels



Metabolomic  
approach



## Next steps in the project


### 1) Drinking water

- Study of the contamination profiles to identify contamination sources
- Study of PFAS contamination in groundwaters (passive sampling)
- Establishing a local fingerprint of PFAS contamination in waters




### 2) Lichens

- Validation of the method
- Analysis of samples from a sampling campaign in Fos-Berre and Lyon



### 3) Toxicity

- Study of the effect of higher concentrations on target genes
- Ecotoxicological assays on aquatic organisms



### 4) Multi-matrix tranfers

- Study of PFAS contamination in soils, plants and groundwater collected in Lyon
- Implementation of a similar campaign in Fos to compare production and usage sites



# Thank you for listening

This work was financially supported by the A\*midex Foundation and by the ERG foundation

Contacts : - chemistry : [maelys.dijoux@etu.univ-amu.fr](mailto:maelys.dijoux@etu.univ-amu.fr)  
- toxicology : [melissa.denton@imbe.fr](mailto:melissa.denton@imbe.fr)

# Bibliography



- [1] 3M, « The science of organic fluorochemistry », *US EPA Administrative Record (AR)* 226-0547, 1999.
- [2] S. E. Fenton *et al.*, « Per- and Polyfluoroalkyl Substance Toxicity and Human Health Review: Current State of Knowledge and Strategies for Informing Future Research », *Environmental Toxicology and Chemistry*, vol. 40, n° 3, p. 606-630, 2021, doi: 10.1002/etc.4890.
- [3] E. P. on C. in the F. C. (EFSA C. Panel) *et al.*, « Risk to human health related to the presence of perfluoroalkyl substances in food », *EFSA Journal*, vol. 18, n° 9, p. e06223, 2020, doi: 10.2903/j.efsa.2020.6223.
- [4] S. Zahm *et al.*, « Carcinogenicity of perfluorooctanoic acid and perfluorooctanesulfonic acid », *The Lancet Oncology*, vol. 25, n° 1, p. 16-17, janv. 2024, doi: 10.1016/S1470-2045(23)00622-8.
- [5] European Parliament, *Directive (EU) 2020/2184 of the European Parliament and of the Council of 16 December 2020 on the quality of water intended for human consumption*. 2020. Consulté le: 31 août 2023. [En ligne]. Disponible sur: <https://eur-lex.europa.eu/eli/dir/2020/2184/oj>
- [6] « Plan d'action ministériel sur les PFAS ». Consulté le: 2 juin 2025. [En ligne]. Disponible sur: <https://www.ecologie.gouv.fr/politiques-publiques/plan-daction-interministeriel-pfas>
- [7] « Arrêté du 31 octobre 2024 relatif à l'analyse des substances per- et polyfluoroalkylées dans les émissions atmosphériques des installations d'incinération, de co-incinération et d'autres traitements thermiques de déchets - Légifrance ». Consulté le: 1 juin 2025. [En ligne]. Disponible sur: <https://www.legifrance.gouv.fr/jorf/id/JORFTEXT000050479313>
- [8] « Installations classées | Géorisques ». Consulté le: 28 mai 2024. [En ligne]. Disponible sur: <https://www.georisques.gouv.fr/risques/installations/donnees/carte#/admin/dpt/13>



# Bibliography



- [9] C. E. Müller et al., « Biomagnification of Perfluorinated Compounds in a Remote Terrestrial Food Chain: Lichen–Caribou–Wolf », *Environ. Sci. Technol.*, vol. 45, no 20, p. 8665-8673, oct. 2011, doi: 10.1021/es201353v.
- [10] J. J. Alava et al., *Perfluorinated Chemicals in Sediments, Lichens, and Seabirds from the Antarctic Peninsula — Environmental Assessment and Management Perspectives*. IntechOpen, 2015. doi: 10.5772/60205.
- [11] W. Nassazzi *et al.*, « A novel method for extraction, clean-up and analysis of per- and polyfluoroalkyl substances (PFAS) in different plant matrices using LC-MS/MS », *Journal of Chromatography B*, vol. 1212, p. 123514, déc. 2022, doi: 10.1016/j.jchromb.2022.123514.
- [12] C. Zhang *et al.*, « Stability of Per- and Polyfluoroalkyl Substances in Solvents Relevant to Environmental and Toxicological Analysis », *Environ. Sci. Technol.*, vol. 56, n° 10, p. 6103-6112, mai 2022, doi: 10.1021/acs.est.1c03979.
- [13] M. Mancini *et al.*, « Evaluation of Pure PFAS Decrease in Controlled Settings », *ACS Meas. Sci. Au*, sept. 2023, doi: 10.1021/acsmeasuresciau.3c00027.
- [14] HCSP Haut conseil de la santé publique, institution française (19/02/2025)
- [15] Mathieu-Denoncourt J, Wallace SJ, de Solla SR, Langlois VS (2015) Plasticizer endocrine disruption: Highlighting developmental and reproductive effects in mammals and non-mammalian aquatic species. *Gen Comp Endocrinol* 219:74–88. <https://doi.org/10.1016/j.ygcen.2014.11.003>
- [16] Cera A, Cesarin G, Spani F, Scalici M (2020) *Hydra vulgaris* assay as environmental assessment tool for ecotoxicology in freshwaters: a review. *Mar Freshw Res* 72:745–753. <https://doi.org/10.1071/MF20115>